arXiv:1604.07289v3 [math.RA] 25 May 2016

Some Identities involving Two Sets of Basis Vectors and the Metric and Mixed Matrices

M.J. Kronenburg

Abstract

Given two sets of basis vectors in n-dimensional space, there exists a relation between their lengths and mutual angles, expressed as relations between the two metric matrices and the mixed matrix. In this paper these relations are given, and their consequences for 2-dimensional and 3-dimensional space are investigated, using a computer algebra program for simplifying expressions.

Keywords: basis vectors, metric matrix, mixed matrix.
MSC 2010: 15A24

1 Introduction

The two metric matrices and the mixed matrix describe the lengths of and mutual angles between two sets of basis vectors in n-dimensional space [5]. There are relations between the two metric matrices and the mixed matrix which are valid for any two sets of basis vectors [4, 5]. These relations can be used in a computer algebra program to find the general identities of the mutual vector angles in 2-dimensional and 3-dimensional space. From these general identities, two special cases [5], one where one of the metric matrices is the identity matrix, and one where the mixed matrix is the identity matrix, can be derived.

2 The Metric and Mixed Matrices in n-dimensional Space

Let $i = 1 \cdots n$ be an index over the dimensions in n-dimensional space, and let $\{\vec{e}_i\}$ be an orthonormal basis in n-dimensional space, which means that:

$$\vec{e}_i \cdot \vec{e}_j = \delta_{ij} \tag{2.1}$$

where δ_{ij} is the Kronecker delta:

$$\delta_{ij} = \begin{cases} 1 & \text{if } i = j \\ 0 & \text{if } i \neq j \end{cases} \tag{2.2}$$

Let the matrices $A = \{A_{ij}\}$ and $A^* = \{A^*_{ij}\}$ have as columns the coordinates on this orthonormal basis of two sets of basis vectors $\{\vec{a}_i\}$ and $\{\vec{a}^*_i\}$, so that [6]:

$$A_{ij} = \vec{e}_i \cdot \vec{a}_j \tag{2.3}$$

$$A^*_{ij} = \vec{e}_i \cdot \vec{a}^*_j \tag{2.4}$$

Let the vector $\vec{r}$ have coordinates on the orthonormal basis $\{\vec{e}_i\}$ and let the vectors $\vec{x}$ and $\vec{x}^*$ be the same as vector $\vec{r}$, but with coordinates on the basis $\{\vec{a}_i\}$ and $\{\vec{a}^*_i\}$, respectively. Then:

$$\vec{r} = A \cdot \vec{x} = A^* \cdot \vec{x}^* \tag{2.5}$$

From this equation follows:

$$\vec{x}^* = A^{*-1} \cdot A \cdot \vec{x} \tag{2.6}$$

Let the lengths of and the mutual angles between the $\{\vec{a}_i\}$ and the $\{\vec{a}^*_i\}$ be given, that is the following matrices are given:

$$G = A^T \cdot A \tag{2.7}$$

$$G^* = A^{*T} \cdot A^* \tag{2.8}$$

$$Q = A^T \cdot A^* \tag{2.9}$$

The matrices G and G^* are the metric matrices [2, 5] and the matrix Q is the mixed matrix [5]. Their elements are $G_{ij} = \vec{a}_i \cdot \vec{a}_j$, $G^*_{ij} = \vec{a}^*_i \cdot \vec{a}^*_j$ and $Q_{ij} = \vec{a}_i \cdot \vec{a}^*_j$. G and G^* are symmetric, and when G (or G^*) is given, a matrix A (or A^*) can be computed with Cholesky decomposition [3]. Furthermore:

$$\vec{r} \cdot \vec{r} = \vec{x} \cdot G \cdot \vec{x} = \vec{x}^* \cdot G^* \cdot \vec{x}^* = \vec{x} \cdot Q \cdot \vec{x}^* = \vec{x}^* \cdot Q^T \cdot \vec{x} \tag{2.10}$$

From (2.9) follows:

$$A^* = (A^{-1})^T \cdot Q \tag{2.11}$$

and equivalently:

$$A = (A^{*-1})^T \cdot Q^T \tag{2.12}$$

Now (2.6) can also be written as [5]:

$$\vec{x}^* = Q^{-1} \cdot G \cdot \vec{x} \tag{2.13}$$

and as

$$\vec{x}^* = G^{*-1} \cdot Q^T \cdot \vec{x} \tag{2.14}$$

Combination of these two identities yields [4]:

$$G^* = Q^T \cdot G^{-1} \cdot Q \tag{2.15}$$

and equivalently

$$G = Q \cdot G^{*-1} \cdot Q^T \tag{2.16}$$

These last two identities can be used to derive some general identities involving the mutual angles between the basis vectors $\{\vec{a}_i\}$ and $\{\vec{a}^*_i\}$ in 2-dimensional and 3-dimensional space.
In the following, when $\cos(\phi) = x$ is known, because ϕ is an angle between two vectors, we can take $0 \leq \phi \leq \pi$, and $\phi = \arccos(x)$.

3 The 2-dimensional case

In the 2-dimensional case, let α_{12} be the angle between $\vec{a}_1$ and $\vec{a}_2$, and β_{12} be the angle between $\vec{a}_1^*$ and $\vec{a}_2^*$, and let γ_{ij} be the angle between $\vec{a}_i$ and $\vec{a}_j^*$. Then the metric and mixed matrices are:

$$G = \begin{pmatrix} |\vec{a}_1|^2 & |\vec{a}_1||\vec{a}_2|\cos(\alpha_{12}) \\ |\vec{a}_1||\vec{a}_2|\cos(\alpha_{12}) & |\vec{a}_2|^2 \end{pmatrix} \tag{3.1}$$

$$G^* = \begin{pmatrix} |\vec{a}_1^*|^2 & |\vec{a}_1^*||\vec{a}_2^*|\cos(\beta_{12}) \\ |\vec{a}_1^*||\vec{a}_2^*|\cos(\beta_{12}) & |\vec{a}_2^*|^2 \end{pmatrix} \tag{3.2}$$

$$Q = \begin{pmatrix} |\vec{a}_1||\vec{a}_1^*|\cos(\gamma_{11}) & |\vec{a}_1||\vec{a}_2^*|\cos(\gamma_{12}) \\ |\vec{a}_2||\vec{a}_1^*|\cos(\gamma_{21}) & |\vec{a}_2||\vec{a}_2^*|\cos(\gamma_{22}) \end{pmatrix} \tag{3.3}$$

By using a computer algebra program, $G^* = Q^T \cdot G^{-1} \cdot Q$ can be evaluated (see below), resulting in:

$$\cos^2(\gamma_{11}) + \cos^2(\gamma_{21}) - 2\cos(\alpha_{12})\cos(\gamma_{11})\cos(\gamma_{21}) = \sin^2(\alpha_{12}) \tag{3.4}$$

$$\cos^2(\gamma_{12}) + \cos^2(\gamma_{22}) - 2\cos(\alpha_{12})\cos(\gamma_{12})\cos(\gamma_{22}) = \sin^2(\alpha_{12}) \tag{3.5}$$

$$\begin{aligned} \cos(\beta_{12}) = \frac{1}{\sin^2(\alpha_{12})}[&\cos(\gamma_{11})\cos(\gamma_{12}) + \cos(\gamma_{21})\cos(\gamma_{22}) \\ &- \cos(\alpha_{12})(\cos(\gamma_{11})\cos(\gamma_{22}) + \cos(\gamma_{12})\cos(\gamma_{21}))] \end{aligned} \tag{3.6}$$

In this case $\sin(\alpha_{12})$ is the area of a parallelogram with side lengths 1 and with α_{12} the mutual angle between the sides.
Subtracting (3.5) from (3.4) results in:

$$\cos(\alpha_{12}) = \frac{\cos^2(\gamma_{11}) + \cos^2(\gamma_{21}) - \cos^2(\gamma_{12}) - \cos^2(\gamma_{22})}{2(\cos(\gamma_{11})\cos(\gamma_{21}) - \cos(\gamma_{12})\cos(\gamma_{22}))} \tag{3.7}$$

This equation can only be used when the denominator is not zero. When this denominator is zero, it follows from (3.4) and (3.5) that the numerator is also zero, so that (3.4) and (3.5) are identical. Using $\sin^2(\alpha_{12}) = 1 - \cos^2(\alpha_{12})$, this equation is quadratic in $\cos(\alpha_{12})$, and solving this quadratic equation yields that in this case $\cos(\alpha_{12}) = \cos(\gamma_{11} \pm \gamma_{21}) = \cos(\gamma_{22} \pm \gamma_{12})$, where the correct sign must be chosen. When this denominator is not zero, and when the four γ_{ij} are given, α_{12} and β_{12} can be computed with the last two identities.

4 The 3-dimensional case

In the 3-dimensional case, let α_{ij} be the angle between $\vec{a}_i$ and $\vec{a}_j$, and β_{ij} be the angle between $\vec{a}_i^*$ and $\vec{a}_j^*$, and let γ_{ij} be the angle between $\vec{a}_i$ and $\vec{a}_j^*$. Then the metric and mixed matrices are:

$$G = \begin{pmatrix} |\vec{a}_1|^2 & |\vec{a}_1||\vec{a}_2|\cos(\alpha_{12}) & |\vec{a}_1||\vec{a}_3|\cos(\alpha_{13}) \\ |\vec{a}_1||\vec{a}_2|\cos(\alpha_{12}) & |\vec{a}_2|^2 & |\vec{a}_2||\vec{a}_3|\cos(\alpha_{23}) \\ |\vec{a}_1||\vec{a}_3|\cos(\alpha_{13}) & |\vec{a}_2||\vec{a}_3|\cos(\alpha_{23}) & |\vec{a}_3|^2 \end{pmatrix} \tag{4.1}$$

$$G^* = \begin{pmatrix} |\vec{a}_1^*|^2 & |\vec{a}_1^*||\vec{a}_2^*|\cos(\beta_{12}) & |\vec{a}_1^*||\vec{a}_3^*|\cos(\beta_{13}) \\ |\vec{a}_1^*||\vec{a}_2^*|\cos(\beta_{12}) & |\vec{a}_2^*|^2 & |\vec{a}_2^*||\vec{a}_3^*|\cos(\beta_{23}) \\ |\vec{a}_1^*||\vec{a}_3^*|\cos(\beta_{13}) & |\vec{a}_2^*||\vec{a}_3^*|\cos(\beta_{23}) & |\vec{a}_3^*|^2 \end{pmatrix} \tag{4.2}$$

$$Q = \begin{pmatrix} |\vec{a}_1||\vec{a}_1^*|\cos(\gamma_{11}) & |\vec{a}_1||\vec{a}_2^*|\cos(\gamma_{12}) & |\vec{a}_1||\vec{a}_3^*|\cos(\gamma_{13}) \\ |\vec{a}_2||\vec{a}_1^*|\cos(\gamma_{21}) & |\vec{a}_2||\vec{a}_2^*|\cos(\gamma_{22}) & |\vec{a}_2||\vec{a}_3^*|\cos(\gamma_{23}) \\ |\vec{a}_3||\vec{a}_1^*|\cos(\gamma_{31}) & |\vec{a}_3||\vec{a}_2^*|\cos(\gamma_{32}) & |\vec{a}_3||\vec{a}_3^*|\cos(\gamma_{33}) \end{pmatrix} \tag{4.3}$$

By using a computer algebra program, $G^* = Q^T \cdot G^{-1} \cdot Q$ can be evaluated (see below), resulting in:

$$\Delta = 2\cos(\alpha_{12})\cos(\alpha_{13})\cos(\alpha_{23}) - \cos^2(\alpha_{12}) - \cos^2(\alpha_{13}) - \cos^2(\alpha_{23}) + 1 \tag{4.4}$$

$$\Omega_1 = \cos(\alpha_{12})\cos(\alpha_{13}) - \cos(\alpha_{23}) \tag{4.5}$$

$$\Omega_2 = \cos(\alpha_{12})\cos(\alpha_{23}) - \cos(\alpha_{13}) \tag{4.6}$$

$$\Omega_3 = \cos(\alpha_{13})\cos(\alpha_{23}) - \cos(\alpha_{12}) \tag{4.7}$$

$$\begin{aligned} &2[\cos(\gamma_{21})\cos(\gamma_{31})\Omega_1 + \cos(\gamma_{11})\cos(\gamma_{31})\Omega_2 + \cos(\gamma_{11})\cos(\gamma_{21})\Omega_3] \\ &+ \cos^2(\gamma_{31})\sin^2(\alpha_{12}) + \cos^2(\gamma_{21})\sin^2(\alpha_{13}) + \cos^2(\gamma_{11})\sin^2(\alpha_{23}) = \Delta \end{aligned} \tag{4.8}$$

$$\begin{aligned} &2[\cos(\gamma_{22})\cos(\gamma_{32})\Omega_1 + \cos(\gamma_{12})\cos(\gamma_{32})\Omega_2 + \cos(\gamma_{12})\cos(\gamma_{22})\Omega_3] \\ &+ \cos^2(\gamma_{32})\sin^2(\alpha_{12}) + \cos^2(\gamma_{22})\sin^2(\alpha_{13}) + \cos^2(\gamma_{12})\sin^2(\alpha_{23}) = \Delta \end{aligned} \tag{4.9}$$

$$\begin{aligned} &2[\cos(\gamma_{23})\cos(\gamma_{33})\Omega_1 + \cos(\gamma_{13})\cos(\gamma_{33})\Omega_2 + \cos(\gamma_{13})\cos(\gamma_{23})\Omega_3] \\ &+ \cos^2(\gamma_{33})\sin^2(\alpha_{12}) + \cos^2(\gamma_{23})\sin^2(\alpha_{13}) + \cos^2(\gamma_{13})\sin^2(\alpha_{23}) = \Delta \end{aligned} \tag{4.10}$$

$$\begin{aligned} \cos(\beta_{12}) = \frac{1}{\Delta}[&(\cos(\gamma_{21})\cos(\gamma_{32}) + \cos(\gamma_{22})\cos(\gamma_{31}))\Omega_1 \\ &+(\cos(\gamma_{11})\cos(\gamma_{32}) + \cos(\gamma_{12})\cos(\gamma_{31}))\Omega_2 \\ &+(\cos(\gamma_{11})\cos(\gamma_{22}) + \cos(\gamma_{12})\cos(\gamma_{21}))\Omega_3 \\ &+\cos(\gamma_{31})\cos(\gamma_{32})\sin^2(\alpha_{12}) + \cos(\gamma_{21})\cos(\gamma_{22})\sin^2(\alpha_{13}) \\ &+\cos(\gamma_{11})\cos(\gamma_{12})\sin^2(\alpha_{23})] \end{aligned} \tag{4.11}$$

$$\begin{aligned} \cos(\beta_{13}) = \frac{1}{\Delta}[&(\cos(\gamma_{21})\cos(\gamma_{33}) + \cos(\gamma_{23})\cos(\gamma_{31}))\Omega_1 \\ &+(\cos(\gamma_{11})\cos(\gamma_{33}) + \cos(\gamma_{13})\cos(\gamma_{31}))\Omega_2 \\ &+(\cos(\gamma_{11})\cos(\gamma_{23}) + \cos(\gamma_{13})\cos(\gamma_{21}))\Omega_3 \\ &+\cos(\gamma_{31})\cos(\gamma_{33})\sin^2(\alpha_{12}) + \cos(\gamma_{21})\cos(\gamma_{23})\sin^2(\alpha_{13}) \\ &+\cos(\gamma_{11})\cos(\gamma_{13})\sin^2(\alpha_{23})] \end{aligned} \tag{4.12}$$

$$\begin{aligned} \cos(\beta_{23}) = \frac{1}{\Delta}[&(\cos(\gamma_{22})\cos(\gamma_{33}) + \cos(\gamma_{23})\cos(\gamma_{32}))\Omega_1 \\ &+(\cos(\gamma_{12})\cos(\gamma_{33}) + \cos(\gamma_{13})\cos(\gamma_{32}))\Omega_2 \\ &+(\cos(\gamma_{12})\cos(\gamma_{23}) + \cos(\gamma_{13})\cos(\gamma_{22}))\Omega_3 \\ &+\cos(\gamma_{32})\cos(\gamma_{33})\sin^2(\alpha_{12}) + \cos(\gamma_{22})\cos(\gamma_{23})\sin^2(\alpha_{13}) \\ &+\cos(\gamma_{12})\cos(\gamma_{13})\sin^2(\alpha_{23})] \end{aligned} \tag{4.13}$$

In this case $\sqrt{\Delta}$ is the volume of a parallelepiped with edge lengths 1 and with the α_{ij} the mutual angles between the edges.

5 The Special Case G=I

When $G = I$, or in other words $\vec{a}_i \cdot \vec{a}_j = \delta_{ij}$, then the basis vectors $\{\vec{a}_i\}$ are an orthonormal basis [5], and (2.15) becomes $G^* = Q^T \cdot Q$. Because $\alpha_{ij} = \pi/2$, $\cos(\alpha_{ij}) = 0$ and $\sin(\alpha_{ij}) = 1$.

5.1 The 2-dimensional case

In the 2-dimensional case, from (3.4), (3.5) and (3.6) follows:

$$\cos^2(\gamma_{11}) + \cos^2(\gamma_{21}) = 1 \tag{5.1}$$

$$\cos^2(\gamma_{12}) + \cos^2(\gamma_{22}) = 1 \tag{5.2}$$

$$\cos(\beta_{12}) = \cos(\gamma_{11})\cos(\gamma_{12}) + \cos(\gamma_{21})\cos(\gamma_{22}) \tag{5.3}$$

Identities (5.1) and (5.2) also follow from trigonometry. Using $\cos^2(x) = 1 - \sin^2(x)$ and $\cos^2(x) - \sin^2(y) = \cos(x+y)\cos(x-y)$ [1], these two identities become:

$$\cos(\gamma_{11} + \gamma_{21})\cos(\gamma_{11} - \gamma_{21}) = 0 \tag{5.4}$$

$$\cos(\gamma_{12} + \gamma_{22})\cos(\gamma_{12} - \gamma_{22}) = 0 \tag{5.5}$$

Because $\alpha_{12} = \pi/2$, there is some sign combination for which:

$$\pm\,\gamma_{11} \pm \gamma_{21} = \frac{\pi}{2} \bmod \pi \tag{5.6}$$

$$\pm\,\gamma_{12} \pm \gamma_{22} = \frac{\pi}{2} \bmod \pi \tag{5.7}$$

and because $\cos(\pi/2 \bmod \pi) = 0$ the two identities follow.
Identity (5.3) also follows from trigonometry.
Using $\cos(x)\cos(y) = (\cos(x+y) + \cos(x-y))/2$ [1] this identity becomes:

$$\cos(\beta_{12}) = \frac{1}{2}[\cos(\gamma_{11} + \gamma_{12}) + \cos(\gamma_{11} - \gamma_{12}) + \cos(\gamma_{22} + \gamma_{21}) + \cos(\gamma_{22} - \gamma_{21})] \tag{5.8}$$

There is some sign combination for which:

$$\pm\,\gamma_{11} \pm \gamma_{12} = \beta_{12} \bmod 2\pi \tag{5.9}$$

$$\pm\,\gamma_{21} \pm \gamma_{22} = \beta_{12} \bmod 2\pi \tag{5.10}$$

The other two terms are $\cos(x) + \cos(y) = 2\cos((x+y)/2)\cos((x-y)/2)$ [1], and one of these arguments becomes $\pi/2 \bmod \pi$.

5.2 The 3-dimensional case

In the 3-dimensional case, because $\Delta = 1$, $\Omega_i = 0$ and $\sin(\alpha_{ij}) = 1$, it follows from (4.8), (4.9) and (4.10):

$$\cos^2(\gamma_{11}) + \cos^2(\gamma_{21}) + \cos^2(\gamma_{31}) = 1 \tag{5.11}$$

$$\cos^2(\gamma_{12}) + \cos^2(\gamma_{22}) + \cos^2(\gamma_{32}) = 1 \tag{5.12}$$

$$\cos^2(\gamma_{13}) + \cos^2(\gamma_{23}) + \cos^2(\gamma_{33}) = 1 \tag{5.13}$$

and it follows from (4.11), (4.12) and (4.13):

$$\cos(\beta_{12}) = \cos(\gamma_{11})\cos(\gamma_{12}) + \cos(\gamma_{21})\cos(\gamma_{22}) + \cos(\gamma_{31})\cos(\gamma_{32}) \tag{5.14}$$

$$\cos(\beta_{13}) = \cos(\gamma_{11})\cos(\gamma_{13}) + \cos(\gamma_{21})\cos(\gamma_{23}) + \cos(\gamma_{31})\cos(\gamma_{33}) \tag{5.15}$$

$$\cos(\beta_{23}) = \cos(\gamma_{12})\cos(\gamma_{13}) + \cos(\gamma_{22})\cos(\gamma_{23}) + \cos(\gamma_{32})\cos(\gamma_{33}) \tag{5.16}$$

6 The Special Case Q=I

When $Q = I$, or in other words $\vec{a}_i \cdot \vec{a}_j^* = \delta_{ij}$, then the basis vectors $\{\vec{a}_i\}$ and $\{\vec{a}_i^*\}$ are mutually reciprocal [5], and (2.15) becomes $G^* = G^{-1}$. Then it follows from (2.11) and (2.12):

$$A^* = (A^{-1})^T \tag{6.1}$$

and equivalently

$$A = (A^{*-1})^T \tag{6.2}$$

Furthermore from (2.10):

$$\vec{r} \cdot \vec{r} = \vec{x} \cdot \vec{x}^* \tag{6.3}$$

From the definition of Q it also follows in this case that the only $\cos(\gamma_{ij})$ unequal to zero are when $i = j$, and

$$|\vec{a}_i||\vec{a}_i^*|\cos(\gamma_{ii}) = 1 \tag{6.4}$$

from which follows that:

$$\cos(\gamma_{ii}) > 0 \tag{6.5}$$

and by taking $0 < \alpha_{ij} < \pi$ we also have:

$$\sin(\alpha_{ij}) > 0 \tag{6.6}$$

6.1 The 2-dimensional case

In the 2-dimensional case, (3.4) and (3.5) with (6.5) and (6.6) give:

$$\cos(\gamma_{11}) = \sin(\alpha_{12}) \tag{6.7}$$

$$\cos(\gamma_{22}) = \sin(\alpha_{12}) \tag{6.8}$$

Combination with (6.4) yields:

$$|\vec{a}_1^*| = \frac{1}{|\vec{a}_1| \sin(\alpha_{12})} \tag{6.9}$$

$$|\vec{a}_2^*| = \frac{1}{|\vec{a}_2| \sin(\alpha_{12})} \tag{6.10}$$

and with (3.6):

$$\cos(\beta_{12}) = -\cos(\alpha_{12}) \tag{6.11}$$

6.2 The 3-dimensional case

In the 3-dimensional case, (4.8), (4.9) and (4.10) with (6.5) and (6.6) give:

$$\cos(\gamma_{11}) \sin(\alpha_{23}) = \sqrt{\Delta} \tag{6.12}$$

$$\cos(\gamma_{22}) \sin(\alpha_{13}) = \sqrt{\Delta} \tag{6.13}$$

$$\cos(\gamma_{33}) \sin(\alpha_{12}) = \sqrt{\Delta} \tag{6.14}$$

Combination with (6.4) yields:

$$|\vec{a}_1^*| = \frac{\sin(\alpha_{23})}{|\vec{a}_1|\sqrt{\Delta}} \tag{6.15}$$

$$|\vec{a}_2^*| = \frac{\sin(\alpha_{13})}{|\vec{a}_2|\sqrt{\Delta}} \tag{6.16}$$

$$|\vec{a}_3^*| = \frac{\sin(\alpha_{12})}{|\vec{a}_3|\sqrt{\Delta}} \tag{6.17}$$

and with (4.11), (4.12) and (4.13):

$$\cos(\beta_{12}) = \frac{\cos(\alpha_{13})\cos(\alpha_{23}) - \cos(\alpha_{12})}{\sin(\alpha_{13})\sin(\alpha_{23})} \tag{6.18}$$

$$\cos(\beta_{13}) = \frac{\cos(\alpha_{12})\cos(\alpha_{23}) - \cos(\alpha_{13})}{\sin(\alpha_{12})\sin(\alpha_{23})} \tag{6.19}$$

$$\cos(\beta_{23}) = \frac{\cos(\alpha_{12})\cos(\alpha_{13}) - \cos(\alpha_{23})}{\sin(\alpha_{12})\sin(\alpha_{13})} \tag{6.20}$$

7 The Inverse Identities

The matrix identity (2.15), that is $G^* = Q^T \cdot G^{-1} \cdot Q$, was used for deriving the above identities. When using matrix identity (2.16), that is $G = Q \cdot G^{*-1} \cdot Q^T$, then the resulting identities are obtained from the above identities by replacing $|\vec{a}_i|$ by $|\vec{a}_i^*|$, $|\vec{a}_i^*|$ by $|\vec{a}_i|$, α_{ij} by β_{ij}, β_{ij} by α_{ij}, and γ_{ij} by γ_{ji}.

8 Computer Program

The Mathematica® [7] programs used to compute the expressions are given below. The 2-dimensional case:

```
G:={{r1^2,r1 r2 Cos[alpha12]},
    {r1 r2 Cos[alpha12],r2^2}}
Q:={{r1 s1 Cos[gamma11],r1 s2 Cos[gamma12]},
    {r2 s1 Cos[gamma21],r2 s2 Cos[gamma22]}}
FullSimplify[Transpose[Q].Inverse[G].Q]
```

The 3-dimensional case:

```
G:={{r1^2,r1 r2 Cos[alpha12],r1 r3 Cos[alpha13]},
    {r1 r2 Cos[alpha12],r2^2,r2 r3 Cos[alpha23]},
    {r1 r3 Cos[alpha13],r2 r3 Cos[alpha23],r3^2}}
Q:={{r1 s1 Cos[gamma11],r1 s2 Cos[gamma12],r1 s3 Cos[gamma13]},
    {r2 s1 Cos[gamma21],r2 s2 Cos[gamma22],r2 s3 Cos[gamma23]},
    {r3 s1 Cos[gamma31],r3 s2 Cos[gamma32],r3 s3 Cos[gamma33]}}
FullSimplify[Transpose[Q].Inverse[G].Q]
```

References

[1] M. Abramowitz, I.A. Stegun, *Handbook of Mathematical Functions*, Dover, 1972.

[2] C. Giacovazzo, H.L. Monaco, G. Artioli, D. Viterbo, G. Ferraris, G. Gilli, G. Zanotti, M. Catti, *Fundamentals of Crystallography*, Oxford University Press, 1992.

[3] G.H. Golub, C.F. van Loan, *Matrix Computations*, Johns Hopkins University Press, 1983.

[4] M.J. Kronenburg, Atomic displacement parameters and anisotropic thermal ellipsoid lengths and angles, *Acta Cryst.* A60 (2004) 250-256.

[5] U. Shmueli, *International Tables for Crystallography, Volume B*, ch. 1.1, Kluwer Academic Publishers, 1993.

[6] K.N. Trueblood, H.B. Bürgi, H. Burzlaff, J.D. Dunitz, C.M. Gramaccioli, H.H. Schulz, U. Shmueli, S.C. Abrahams, Atomic Displacement Parameter Nomenclature, *Acta Cryst.* A52 (1996) 770-781.

[7] S. Wolfram, *The Mathematica Book*, 5th ed., Wolfram Media, 2003.